



## When your child has been diagnosed with ADHD



### Choose: Consistency

Discover medication that works with your child's unique profile for consistent, predictable symptom control

#### IMPORTANT SAFETY INFORMATION

What is the most important information I should know about Quillivant XR® (methylphenidate HCl) and QuilliChew ER® (methylphenidate HCl)?

**Quillivant XR and QuilliChew ER are federally controlled substance (CII) because they can be a target for people who abuse prescription medicines or street drugs.** Quillivant XR and QuilliChew ER has a high chance for abuse and misuse and may lead to substance use problems, including addiction. **Keep the medicines in a safe place to prevent abuse and misuse.** Never give your Quillivant XR or QuilliChew ER to anyone else, because it may cause death or harm them. **Selling or giving away these medicines may harm others and is against the law.**

Please see additional Important Safety Information throughout, and [Full Prescribing Information](#), including **Boxed Warning about Abuse, Misuse and Addiction**, and [Medication Guide](#).



## About ADHD

**ADHD is one of the most common neurodevelopmental disorders of childhood<sup>1</sup>. According to a 2022 study in the US, among children aged 3-17:**

**~11%** have ever had a diagnosis of ADHD<sup>2</sup>

**~54%** of those with current ADHD take medication to manage symptoms<sup>2</sup>

But finding the right medication can be difficult. A recent study found:

**~60%** of children and adolescents changed ADHD treatments over a 12-month period<sup>3</sup>

As the parent of a child with ADHD, you know how important consistency is to their daily routine.<sup>4</sup>

Learn more in this brochure about **Quillivant XR** (methylphenidate HCl) and **QuilliChew ER** (methylphenidate HCl) — two ADHD medications that work with your child's unique profile to provide consistent symptom control you can count on.

### IMPORTANT SAFETY INFORMATION (cont'd)

**Your healthcare provider should check you or your child's risk for abuse, misuse, and addiction before starting treatment with Quillivant XR or QuilliChew ER and will monitor you or your child during treatment.** Tell your physician if you, your child, or any family members have ever abused or been dependent on alcohol, prescription medicines, or street drugs.

**Quillivant XR and QuilliChew ER should not be taken if you or your child** are allergic to methylphenidate hydrochloride, or any of the ingredients in QuilliChew ER, or are taking or have taken within the past 14 days an antidepressant medicine called a monoamine oxidase inhibitor or MAOI.

**Please see additional Important Safety Information throughout, and [Full Prescribing Information](#), including **Boxed Warning about Abuse, Misuse and Addiction**, and [Medication Guide](#).**



**TWO FORMULATIONS** to meet your child's specific needs.

**ONE UNIQUE TECHNOLOGY** to work with your child's unique profile.



**Continuous** release through the most important parts of the day<sup>5,6,7</sup>



**Consistent**, smooth drug delivery<sup>5</sup>



**Predictable**, long-lasting symptom control<sup>6,7</sup>

### How does our technology work?

A special coating on drug delivery particles **protects** methylphenidate until it's ready to be released - so differences in your child's metabolism or changes in their diet don't affect how the medication is released through the day.<sup>5</sup>

### IMPORTANT SAFETY INFORMATION (cont'd)

The following have been reported with the use of methylphenidate hydrochloride and other stimulant medicines:

• **Heart-related problems:**

- Sudden death in patients who have heart problems or heart defects.
- Stroke and heart attack in adults.
- Increased blood pressure and heart rate.
- Your healthcare provider should check your or your child's blood pressure and heart rate regularly during treatment with Quillivant XR or QuilliChew ER.

Please see additional Important Safety Information throughout, and [Full Prescribing Information](#), including **Boxed Warning about Abuse, Misuse and Addiction**, and [Medication Guide](#).

## Meet Quillivant XR (methylphenidate HCl) oral suspension

### Helps control symptoms from morning through the day<sup>7</sup>

#### Continuous 12-hour symptom control<sup>7</sup>

- Lasting improvement in attention and behavior<sup>7</sup>

#### Consistent Medication Delivery

- Continuous, smooth release of methylphenidate throughout the day<sup>5</sup>
- Release not affected by food consumption or metabolism of methylphenidate<sup>5</sup>

#### IMPORTANT SAFETY INFORMATION (cont'd)

##### • Mental (psychiatric) problems:

- New or worse behavior and thought problems.
- New or worse bipolar illness.
- May cause new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms.

##### • Circulation problems in fingers and toes (peripheral vasculopathy, including Raynaud's phenomenon):

- Fingers or toes may feel numb, cool, painful, or may change color from pale, to blue, to red.

#### Call your healthcare provider right away if you or your child have any:

- Heart-related symptoms such as chest pain, shortness of breath, or fainting.
- New or worsening mental (psychiatric) symptoms or new manic symptoms.
- Signs of unexplained wounds appearing on fingers or toes while taking Quillivant XR or QuilliChew ER.

#### Quillivant XR or QuilliChew ER may not be right for you or your child. Before starting Quillivant XR, tell your or your child's health care provider about all health conditions including:

- Heart problems, heart disease, heart defects, or high blood pressure.
- mental problems including psychosis, mania, bipolar illness, or depression.
- circulation problems in fingers and toes.
- have eye problems, including increased pressure in your eye, glaucoma, or problems with your close-up vision (farsightedness).
- have or had repeated movements or sounds (tics) or Tourette's syndrome, or have a family history of tics or Tourette's syndrome.


Please see additional Important Safety Information throughout, and [Full Prescribing Information](#), including **Boxed Warning about Abuse, Misuse and Addiction**, and [Medication Guide](#).



## Personalized dosing in a long-acting liquid<sup>8</sup>

Liquid Quillivant XR (methylphenidate HCl) allows your doctor to adjust the dose, up or down, weekly, in amounts of 2 to 4 mL.<sup>8</sup>

- Quillivant XR comes with an oral dosing dispenser to help you measure the correct amount of medication
- Quillivant XR can be taken with or without food
- Take Quillivant XR exactly as prescribed and remember that you **MUST** shake the bottle vigorously for at least 10 seconds prior to each use



**Please see Instructions for Use in the accompanying Full Prescribing Information in pocket of this brochure for specific instructions on how your child should take Quillivant XR.**

Or visit [www.trisadhd.com/quillivant-xr/taking-quillivant-xr/](http://www.trisadhd.com/quillivant-xr/taking-quillivant-xr/)

### IMPORTANT SAFETY INFORMATION (cont'd)

**Quillivant XR (methylphenidate HCl) or QuilliChew ER (methylphenidate HCl) may not be right for you or your child. Before starting Quillivant XR, tell your or your child's health care provider about all health conditions including (cont'd):**

- phenylketonuria (PKU). QuilliChew ER extended-release chewable tablets contain phenylalanine as part of the artificial sweetener, aspartame. The artificial sweetener may be harmful to people with PKU or who are allergic to phenylalanine.
- if you are pregnant or plan to become pregnant. It is not known if Quillivant XR or QuilliChew ER will harm the unborn baby. There is a pregnancy registry for females who are exposed to QuilliChew ER during pregnancy. The purpose of the registry is to collect information about the health of females exposed to Quillivant XR or QuilliChew ER and their baby. If you or your child becomes pregnant during treatment with QuilliChew ER, talk to your healthcare provider about registering with the National Pregnancy Registry for Psychostimulants at 1-866-961-2388 or visit <https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/othermedications/>.

**Please see additional Important Safety Information throughout, and [Full Prescribing Information](#), including [Boxed Warning about Abuse, Misuse and Addiction](#), and [Medication Guide](#).**

## Meet QuilliChew ER (methylphenidate HCl) tablets

### When **tablets** are the preferred option for your child

#### Effective control of symptoms<sup>6</sup>

- Works through the most important parts of the day<sup>6</sup>
- Lasting improvements in attention and behavior<sup>6</sup>

#### Consistent Delivery — chewed or swallowed<sup>9</sup>

- Continuous, smooth release of medication<sup>5</sup>
- Release not affected by diet or metabolism<sup>5</sup>

#### IMPORTANT SAFETY INFORMATION (cont'd)

**Quillivant XR (methylphenidate HCl) or QuilliChew ER (methylphenidate HCl) may not be right for you or your child. Before starting Quillivant XR, tell your or your child's health care provider about all health conditions including (cont'd)**

- if you are breastfeeding or plan to breast feed. Quillivant XR or QuilliChew ER passes into your breast milk. You and your healthcare provider should decide if you will take QuilliChew ER or breastfeed.

**Do not start any new medicine while taking Quillivant XR or QuilliChew ER without talking to your healthcare provider first.**

#### What should I avoid while taking Quillivant XR or QuilliChew ER?

Quillivant XR and QuilliChew ER should not be taken with MAOI medicines or if you stopped taking an MAOI in the last 14 days. Do not drink alcohol while taking Quillivant XR or QuilliChew ER. This may cause a faster release of your methylphenidate dose.

#### Quillivant XR or QuilliChew ER may cause serious side effects, including:

- See “What is the most important information I should know about Quillivant XR or QuilliChew ER?” for information on reported heart and mental problems.

#### Other serious side effects include:

- **New or worsening tics or worsening Tourette's syndrome.** Tell your healthcare provider if you or your child get any new or worsening tics or worsening Tourette's syndrome during treatment with **Quillivant XR or QuilliChew ER.**
- **Painful and prolonged erections (priapism)** have occurred with methylphenidate. If you or your child develop priapism seek medical help right away. **Because priapism can cause long lasting damage, it should be checked by a health care provider right away.**

Please see additional Important Safety Information throughout, and [Full Prescribing Information](#), including **Boxed Warning about Abuse, Misuse and Addiction**, and [Medication Guide](#).



## Personalized dosing for your child

QuilliChew ER (methylphenidate HCl) should be taken by mouth once in the morning.

- Can be chewed or swallowed whole<sup>9</sup>
- Can be taken with or without food

## 11 dosing options<sup>10</sup>

- The 20-mg and 30-mg tablets are scored<sup>10</sup>
  - Scored tablets can be divided into equal halves to allow additional dosing options and to achieve the specific dose that is recommended by your healthcare provider<sup>10</sup>
  - The 20-mg tablet can be split into 10-mg equal halves, and the 30-mg tablet can be split into 15-mg equal halves<sup>10</sup>
  - The 40-mg tablet is not scored<sup>10</sup>



Or visit [www.trisadhd.com/quillichew-er/taking-quillichew-er/](http://www.trisadhd.com/quillichew-er/taking-quillichew-er/)

### IMPORTANT SAFETY INFORMATION (cont'd)

#### Other serious side effects include (cont'd):

- **Slowing of growth (height and weight) in children.** Children should have their height and weight checked often during treatment with Quillivant XR (methylphenidate HCl) or QuilliChew ER (methylphenidate HCl). Quillivant XR or QuilliChew ER treatment may be stopped if your child is not gaining weight or height.
- **Eye problems (increased pressure in the eye and glaucoma).** Call your healthcare provider right away if you or your child develop changes in your vision or eye pain, swelling, or redness.

Please see additional Important Safety Information throughout, and [Full Prescribing Information](#), including **Boxed Warning about Abuse, Misuse and Addiction**, and [Medication Guide](#).




## Ready to switch?

### Talk to your doctor about whether your child is ready to transition from a **liquid medication** to a **chewable tablet**.

Both **Quillivant XR** (methylphenidate HCl) and **QuilliChew ER** (methylphenidate HCl) use the same unique delivery technology, so whichever formulation you choose, you can expect:

#### **Consistent Delivery — chewed or swallowed<sup>9</sup>**

- Consistent ADHD medication delivery<sup>5</sup>
- Effective ADHD symptom control<sup>6,7</sup>



#### **IMPORTANT SAFETY INFORMATION (cont'd)**

##### **Common side effects include:**

- |                      |                |                   |                            |
|----------------------|----------------|-------------------|----------------------------|
| - Decreased appetite | - Indigestion  | - Dizziness       | - Increased blood pressure |
| - Trouble sleeping   | - Stomach pain | - Irritability    |                            |
| - Nausea             | - Weight loss  | - Mood swings     |                            |
| - Vomiting           | - Anxiety      | - Fast heart beat |                            |

These are not all the possible side effects of Quillivant XR and QuilliChew ER.

**Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

#### **APPROVED USE**

**Quillivant XR** (methylphenidate HCl) and **QuilliChew ER** (methylphenidate HCl) are central nervous system stimulant prescription medicines. **Quillivant XR is a liquid medicine** that you take by mouth. **QuilliChew ER is a chewable tablet.**

Quillivant XR and QuilliChew ER are used for the treatment of Attention Deficit Hyperactivity Disorder (ADHD). Quillivant XR and QuilliChew ER may help increase attention and decrease impulsiveness and hyperactivity in people with ADHD.

**Quillivant XR, and QuilliChew ER are not recommended for use in children under 6 years of age with ADHD.**

**Please see additional Important Safety Information throughout, and [Full Prescribing Information](#), including Boxed Warning about Abuse, Misuse and Addiction, and [Medication Guide](#).**

**TWO FORMULATIONS** to meet your child's specific needs.

**ONE UNIQUE TECHNOLOGY** to work with your child's unique profile.



**Efficacy** through the most important parts of the day<sup>6,7</sup>



**Lasting** improvement in attention and behavior<sup>6,7</sup>

If you're switching between **Quillivant XR** and **QuilliChew ER**, follow your healthcare provider's step-by-step dosing plan for the new medicine. QuilliChew ER and Quillivant XR doses are not directly interchangeable.



**Join our community!**



**Visit our websites and sign up  
for our newsletter**

[quillivantXR.com](http://quillivantXR.com)

[quillichewER.com](http://quillichewER.com)



**Connect on Facebook**

[Quillivant XR](#)

[QuilliChew ER](#)

Please see additional Important Safety Information throughout, and [Full Prescribing Information](#), including Boxed Warning about Abuse, Misuse and Addiction, and [Medication Guide](#).

For people 6 years and older with Attention Deficit Hyperactivity Disorder (ADHD)



## Consistent ADHD medication delivery<sup>5</sup> Effective ADHD symptom control<sup>6,7</sup>



Eligible patients pay as little  
as **\$25** each month\*

Learn how you can pay as little  
as **\$25** for your prescription at  
[trissavingscard.com](http://trissavingscard.com)

Text TRIS to TRIS4U (874748) for a  
Mobile Savings Card.\*\*

\* Terms and conditions apply. Card issue is subject to eligibility.

\*\* Mobile Terms and Conditions: [TrisSavingsCard.com/terms](http://TrisSavingsCard.com/terms). Message and Data rates may apply

**References:** **1.** CDC. What is ADHD. <https://cdc.gov/adhd/about> **2.** Danielson ML, Claussen AH, Bitsko RH, Katz SM, Newsome K, Blumberg SJ, Kogan MD, Ghandour R. ADHD Prevalence Among U.S. Children and Adolescents in 2022: Diagnosis, Severity, Co-Occurring Disorders, and Treatment. *J Clin Child Adolesc Psychol.* 2024 May-Jun; 53(3):343-360. **3.** Schein J, Childress A, Adams J, Gagnon-Sanschagrin P, Maitland J, Qu W, Cloutier M, Guérin A. Treatment patterns among children and adolescents with attention-deficit/hyperactivity disorder in the United States - a retrospective claims analysis. *BMC Psychiatry.* 2022 Aug 18;22(1):555. **4.** The World Federation of ADHD guide [electronic resource] / Editors, Luis Augusto Rohde... [et al.]. - Porto Alegre : Artmed, 2019. **5.** Cutler AJ, Hanaie J. Understanding the delivery technology used in ADHD stimulant medications can help to individualize treatment. *CNS Spectrums.* 2025;30(1):e30. **6.** Wigal SB, Childress A, Berry SA, et al. Efficacy and safety of a chewable methylphenidate extended-release tablet in children with attention deficit/ hyperactivity disorder. *J Child Adolesc Psychopharmacol.* 2017;27(8):690-699. **7.** Wigal SB, Childress AC, Belden HW, Berry SA. NWP06, an extended-release oral suspension of methylphenidate, improved attention-deficit/hyperactivity disorder symptoms compared with placebo in a laboratory classroom study. *J Child Adolesc Psychopharmacol.* 2013;23(1):3-10. **8.** Quillivant XR [package insert]. Tris Pharma, Inc., Monmouth Junction, NJ. **9.** Abbas R, Childress AC, Nagraj P, Rolke R, Berry SA, Palumbo DR. Relative bioavailability of methylphenidate extended-release chewable tablets chewed versus swallowed whole. *Clin Ther.* 2018;40(5):733-740. **10.** QuilliChew ER [package insert]. Tris Pharma, Inc., Monmouth Junction, NJ.

Please see additional Important Safety Information throughout, and [Full Prescribing Information](#), including **Boxed Warning about Abuse, Misuse and Addiction**, and [Medication Guide](#).



Quillivant XR and QuilliChew ER are registered trademarks of NextWave Pharmaceuticals, Inc.  
© 2026 Tris Pharma, Inc. All rights reserved. QUI.2367.PR 01/26